

# EOS MLS Upper Tropospheric and Lower Stratospheric Ozone Validation by Sonde Measurements & Other Ground-Based Datasets

Yibo Jiang<sup>1</sup>, L. Froidevaux<sup>1</sup>, A. Lambert<sup>1</sup>, N. J. Livesey<sup>1</sup>, W. G. Read<sup>1</sup>, J.W. Waters<sup>1</sup>, B. Bojkov<sup>2</sup>

<sup>1</sup>Jet Propulsion Laboratory, California Institute of Technology

<sup>2</sup> NASA Goddard Space Flight Center


+ other contributors (lidar data, JPL team)



#### Ozone Sonde Site Distribution (used in this analysis)

Available data from AVDC


- Good coverage in tropics and northern hemisphere
- Poor coverage in southern hemisphere, Europe and Asia





#### **Sample Profile Comparisons**

- Coincidence Profiles
  - Same day
  - Location within
    1º latitude
    12º longitude
- Least-squares fit is used here to fit the sonde profiles to the MLS vertical retrieval grid






# Scatter Plots (color coded by latitude)


#### MLS V1.5 & Sondes

- Good correlation except at 316 hPa in equatorial and subtropics region (black, green, red)
- Upper stratospheric sonde data quality probably influences poorer comparisons there






# Globally-averaged comparisons MLS V1.5 & Sondes



 The average differences are fairly constant in time







#### Latitude Distribution (at different pressure levels)

MLS V1.5 & Sondes

- There is generally good tracking versus latitude down to 215 hPa
- The differences are most often within 20% or better down to 215 hPa, except in the tropics








#### Latitude Distribution (Column Ozone for LS)

MLS V1.5 & Sondes


Many differences are within 3%;
 need to look at the larger differences
 more carefully





#### MLS Ozone v2.1, v1.5 & Sondes

- Based on 17 days of MLS v2.1 data
   58 coincidence profiles (available on AVDC)
- MLS v2.1 O3 shows better agreement with sonde data than v1.5
- within 5% from ~ 20 to 100 hPa
- 10% in US, MLS biased lower
- still biased high in UT, but improved over v1.5
- Upper stratospheric sonde data quality probably influences poorer comparisons there
- As for comparisons vs satellite data, we see the change in slope for MLS v2.1



### Comparisons with O3 LIDAR Data




#### Times Series of MLS Ozone v1.5 & LIDAR

- 2 Long-term LIDAR Measurement Sites
  - •Table Mountain(34.5N, 117.7W)
  - •Mauna Loa(19.5N, 155.7W)
- Seasonal Cycles can be seen in both datasets at 15 hPa
- •MLS and LIDAR are tracking each other very well at 100 hPa in Table Mountain even data has large variability


  Mauna Loa

**Table Mountain** 

# Pres=100 hPa



#### Pres=15 hPa




### Comparisons with O3 LIDAR Data



### MLS Ozone v2.1, v1.5 & LIDAR

- Based on 17 days of MLS v2.1 data
   18 coincidence profiles (available on AVDC)
- MLS v2.1 O3 shows better agreement with LIDAR data than v1.5
- v2.1 lidar within 5-10%
   for P between 215 and 0.7 hPa
- MLS sondes differences near 10 hPa is not observed in LIDAR comparison
- MLS is biased high at 315 hPa



#### Aura MLS Ozone UT/LS Validation



### **Summary**

#### MLS V2.1 Data

- V2.1 shows better overall agreement than V1.5 compared to sondes & LIDAR
- The differences between MLS and sondes are within 10%, except at pressure 215 & 316 hPa based on 17 days available MLS v2.1 data
- MLS v2.1 and LIDAR within 5-10% for P between 215 and 0.7 hPa
- There is still a high bias in UT especially in tropics

### **Aura MLS Ozone UT/LS Validation**



# **Summary (cont.)**

#### Validation plans and paper

- Would be good to add more sondes data from European sites
- Some issues with lidar data files (now on AVDC)

#### **Comparisons with others datasets**

 Ground-Based Microwave Data (some work presented by others at Sep. 2005 meeting)

Please discuss any related plans for validation papers (for JGR special issue) with MLS team